

RB Health (US) LLC

-PE Product Labels –

-continued

Drug Facts	Drug Facts (continued)
Active ingredients (in each caplet) Acetaminophen 325 mg... Pain reliever Dextromethorphan HBr 10 mg... Cough suppressant Guaifenesin 200 mg... Expectorant Pseudoephedrine HCl 5 mg... Nasal decongestant	Directions ■ cough that occurs with too much mucus (mucus) Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product do not use more than directed. Stop use and ask a doctor if: ■ nausea, dizziness, or sleepiness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 12 caplets in 24 hours, which is the maximum daily amount. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.	Other information ■ store at 20-25°C (68-77°F) Inactive ingredients croscarmellose sodium, croscopolidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pyridone, talc, titanium dioxide.
Questions? 1-866-MUCINEX (1-866-662-4639) You may also report side effects to this phone number.	

MAXIMUM STRENGTH

Mucinex[®]
SINUS-MAX[®]

PRESSURE, PAIN & COUGH

PARENTS:

 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Maximum Strength per 4-hour dose

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

Keep carton for complete product information.

MAXIMUM STRENGTH

NDC 63824-242-20

Mucinex[®]
SINUS-MAX[®]

Actual Size



PRESSURE, PAIN & COUGH

 Acetaminophen – Pain Reliever • Dextromethorphan HBr – Cough Suppressant
 Guaifenesin – Expectorant • Phenylephrine HCl – Nasal Decongestant

- ✓ Relieves Sinus Pressure & Congestion
- ✓ Relieves Headache
- ✓ Controls Cough
- ✓ Thins & Loosens Mucus

FOR AGES 12+

20 CAPLETS

 HEALTH • HYGIENE • HOME

 Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224
 ©2020 RB Health

 www.mucinex.com
 Patents: www.rb.com/patents
 092520 3165986


3

63824 24220

0

 LOT:
 EXP:
 MADE IN:
 3165986







Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224
©2017 RB

101917

LOT:

EXP:

MADE IN:

3055094

- nasal congestion
- minor aches and pains
- sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- headache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

PEEL HERE

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Please visit our website
www.mucinex.com

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Mucinex[®]
Children's
COLD & FLU

Acetaminophen 325 mg – Pain Reliever/Fever Reducer
Dextromethorphan HBr 10 mg – Cough Suppressant
Guaifenesin 200 mg – Expectorant
Phenylephrine HCl 5 mg – Nasal Decongestant

Pediatrician Recommended[†]

NDC 63824-954-04

Mucinex[®]
Children's
COLD & FLU

Acetaminophen 325 mg – Pain Reliever/Fever Reducer
Dextromethorphan HBr 10 mg – Cough Suppressant
Guaifenesin 200 mg – Expectorant
Phenylephrine HCl 5 mg – Nasal Decongestant

ALL IN ONE^{*}

HEADACHE
BODY PAIN
SORE THROAT
FEVER
CHEST CONGESTION
COUGH
NASAL CONGESTION
SINUS CONGESTION
SINUS PRESSURE

Ages 6+ yrs


Mucinex[®]
Children's
COLD & FLU

Each dose of Very Berry flavored Mucinex Children's Cold & Flu Liquid temporarily relieves:


ALL IN ONE^{*}

HEADACHE
BODY PAIN
SORE THROAT
FEVER
CHEST CONGESTION
COUGH
NASAL CONGESTION
SINUS CONGESTION
SINUS PRESSURE





Very Berry
Flavor Liquid




4 FL OZ (118 mL)

Dosing Cup Included

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health
Made in Mexico




HEALTH • HYGIENE • HOME

021221 3163801 US Patents: www.rb.com/patents



3 63824 27864 3

LOT CODE: [REDACTED]
EXP. DATE: [REDACTED]



1083913 3163801

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg...	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash, or

2937

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)

■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

***as an OTC brand in the Children's Cough/Cold category (AlphaImpactRxProVoice Survey)**

***Helps to relieve these symptoms day or night**

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Keep carton for full information.

3163801



Mucinex
Junior

Tamper evident: Do not use if
carton is damaged or if
printed seal on blister is
broken or missing.

COUGH & CONGESTION

Pediatrician Recommended†

NDC 63824-671-20

NEW

Mucinex
Junior

COUGH & CONGESTION

Dextromethorphan HBr 5 mg – Cough Suppressant
Guaifenesin 100 mg – Expectorant
Phenylephrine HCl 2.5 mg – Nasal Decongestant

Ages **6+** yrs

- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Relieves Stuffy Nose
- ✓ Controls Cough

Junior Caplet®



Actual Size

20 CAPLETS

DO NOT CHEW

LOT & EXP.
DATE



0
63824 60068
3

www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2019 RB Health
031419 3098704



HEALTH • HYGIENE • HOME

US Patents: www.rb.com/patents

B-1715-720-09
ORG
DATA MATRIX

Drug Facts

Active ingredients (in each caplet)

Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 100 mg.....Expectorant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough that lasts such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occurs with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: give 2 caplets every 4 hours while symptoms last; do not give more than 6 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, magnesium stearate, maltodextrin, mica, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

***as an OTC brand in the Children's Cough/Cold category (AlphaImpactRxProVoice Survey)**

***as compared to Mucinex caplets**

PARENTS:


Learn about teen medicine abuse


www.StopMedicineAbuse.org

Keep carton for full information.
If your child has difficulty swallowing solid dosage forms, try our Mucinex Children's liquids or mini-melts products.

17.0000

B-1715-720-09
ORG





COUGH & CONGESTION

Each dose of Very Berry flavored
Mucinex Children's
Cough & Congestion Liquid:

- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Relieves Stuffy Nose
- ✓ Controls Cough



Dosing Cup Included

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2019 RB Health
Made in England
021119 3091844



HEALTH ▸ HYGIENE ▸ HOME

■ stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

[illegible]

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Mucinex[®]
Children's

FREEFROM
MULTI-SYMPTOM
COLD & FLU NIGHTTIME

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT
TRIPROLIDINE HCl – ANTIHISTAMINE

Drug Facts (continued)

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 4 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, cochineal camine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

Pediatrician Recommended[†]

NIGHTTIME

NDC 63824-959-64


Mucinex[®]
Children's

FREEFROM

MULTI-SYMPTOM
COLD & FLU
NIGHTTIME

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT
TRIPROLIDINE HCl – ANTIHISTAMINE





relief you trust - **FREEFROM** the
unwanted additives:

Caffeine Free · No Artificial Colors,
Dyes, or Flavors · Alcohol & Sugar Free
No High Fructose Corn Syrup

POWERFUL SYMPTOM RELIEF

- ✓ Nighttime Relief for a Better Morning
- ✓ Relieves Sore Throat & Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Runny Nose & Sneezing
- ✓ Reduces Fever

DOSING CUP INCLUDED



ELDERBERRY
& CHERRY NATURAL FLAVOR

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

HEALTH • HYGIENE • HOME

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - runny nose
 - cough
 - temporarily reduces fever
 - controls cough to help your child get to sleep
- headache
- nasal congestion
- sneezing
- stuffy nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 1625 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- diabetes
- thyroid disease
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- glaucoma

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Tamper evident. Do not use if neckband on bottle cap is broken or missing.



Mucinex[®] Children's FREEFROM
MULTI-SYMPTOM
COLD & STUFFY NOSE
DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
GUAIFENESIN - EXPECTORANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, edotato disodium, flavors, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

*vs Private Label OTC
as an OTC brand in the Children's Cough/Cold category
(AlphaImpactRxProVoice Survey)

Pediatrician Recommended[†]



DAYTIME

NDC 6328-954-04



FREEFROM

**MULTI-SYMPTOM
COLD
& STUFFY NOSE**

DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
GUAIFENESIN - EXPECTORANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT



FREEFROM
ARTIFICIAL FLAVORS, COLORS, & DYES • ALCOHOL & SUGAR

**HONEY
& BERRY NATURAL FLAVOR**



**AGES
4+ YRS.**

4 FL OZ (118 mL)

What this has is just as important as what it doesn't.

Leading Children's OTC Cough & Cold brand* with the effective symptom relief you trust - FREEFROM the unwanted additives:

Caffeine Free • No Artificial Colors, Dyes, or Flavors • Alcohol & Sugar Free
No High Fructose Corn Syrup

POWERFUL SYMPTOM RELIEF

- ✓ Relieves Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Chest Congestion
- ✓ Thins & Loosens Mucus

DOSING CUP INCLUDED



**HONEY
& BERRY NATURAL FLAVOR**

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Questions or comments?
1-866-Mucinex (1-866-682-4639)
www.mucinex.com

HEALTH • HYGIENE • HOME

Drug Facts

Active ingredients (in each 5 mL)	Purposes
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health

010720 3132568 US Patents: www.rb.com/patents
Keep carton for full information.



3 63824 60069 7

LOT CODE:
EXP. DATE:
MADE IN:

3132568

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Mucinex[®]
Children's

FREEFROM

MULTI-SYMPTOM
COLD, FLU & SORE THROAT

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
GUAIFENESIN – EXPECTORANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use


Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

Pediatrician Recommended†

 **DAYTIME**


NDC 63624-958-64

Mucinex[®]
Children's

FREEFROM

MULTI-SYMPTOM
COLD, FLU & SORE THROAT

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
GUAIFENESIN – EXPECTORANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT



*vs Private Label OTC
†as an OTC brand in the Children's Cough/Cold category



relief you trust – **FREEFROM** the
unwanted additives:

Caffeine Free • No Artificial Colors,
Dyes, or Flavors • Alcohol & Sugar Free
No High Fructose Corn Syrup

POWERFUL SYMPTOM RELIEF

- ✓ Relieves Sore Throat
- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion & Cough
- ✓ Thins & Loosens Mucus
- ✓ Reduces Fever

DOSING CUP INCLUDED



ELDERBERRY
& CHERRY NATURAL FLAVOR

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com



HEALTH • HYGIENE • HOME

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
 - nasal congestion
 - sore throat
 - stuffy nose
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- high blood pressure
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- heart disease
- thyroid disease

Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)



3163772

Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....	Cough suppressant
Guaifenesin 100 mg.....	Expectorant
Phenylephrine HCl 2.5 mg.....	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

*as an OTC brand in the Children's Cough/Cold category (AlphaImpactRxProVoice Survey)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Other convenient forms for Chest Congestion Relief

Mucinex Children's
CHEST CONGESTION

mini-melts™

On-the-go granule packets for chest congestion relief!

Keep carton for full information

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Mucinex Children's
MULTI-SYMPTOM COLD

Dextromethorphan HBr 5 mg - Cough Suppressant
Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

Pediatrician Recommended†

NDC 63824-040-01

Mucinex Children's
MULTI-SYMPTOM COLD

Dextromethorphan HBr 5 mg - Cough Suppressant
Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

- ✓ Relieves Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus



Ages 4+ yrs

Very Berry Flavor Liquid

4 FL OZ (118 mL)

Mucinex Children's
MULTI-SYMPTOM COLD

Each dose of Very Berry flavored Mucinex Children's Multi-Symptom Cold Liquid:

- ✓ Relieves Stuffy Nose
- ✓ Controls Cough
- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus*



Dosing Cup Included

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Dist by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health
Made in Mexico
020421 3163772

rb
HEALTH • HYGIENE • HOME

3163772



3 63824 01064 9

LOT CODE:
EXP. DATE:



Drug Facts (continued)

Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

†as an OTC brand in the Children's Cough/Cold

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Mucinex[®]
Children's

**MULTI-SYMPTOM
COLD & SORE THROAT**

Acetaminophen 325 mg - Pain Reliever/Fever Reducer
Dextromethorphan HBr 10 mg - Cough Suppressant
Guaifenesin 200 mg - Expectorant
Phenylephrine HCl 5 mg - Nasal Decongestant

Pediatrician Recommended[†]

NDC 63824-952-17

Mucinex[®]
Children's

**MULTI-SYMPTOM
COLD &
SORE THROAT**

Acetaminophen 325 mg - Pain Reliever/Fever Reducer
Dextromethorphan HBr 10 mg - Cough Suppressant
Guaifenesin 200 mg - Expectorant
Phenylephrine HCl 5 mg - Nasal Decongestant

- ✓ Relieves Sore Throat
- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion and Cough
- ✓ Breaks up Mucus



Ages **6+** yrs

Very Berry
Flavor Liquid

category (AlphaImpactRxProVoice Survey)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org


Keep carton for full information.

4 FL OZ (118 mL)



LOT CODE:

EXP. DATE:



Mucinex[®]
Children's

**MULTI-SYMPTOM
COLD &
SORE THROAT**

Drug Facts

**Active ingredients
(in each 10 mL)**

Purposes

Acetaminophen 325 mg...	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - minor aches and pains
 - sore throat
 - stuffy nose
 - headache

**Each dose of Very Berry flavored
Mucinex Children's Multi-Symptom
Cold & Sore Throat Liquid:**

- ✓ Relieves Sore Throat
- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion and Cough
- ✓ Breaks up Mucus



Dosing Cup Included

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2018 RB Health
Made in Mexico
110118 3078385



HEALTH • HYGIENE • HOME

- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus) ▶



RB Artwork and Print Specification

Product Reference ID: RB123456
 Date: 10/10/2024
 Version: 1.0
 Category: Children's
 Target Audience: Parents
 Product Name: Mucinex Children's
 Product Description: Multi-Symptom Cold
 Product Type: Liquid
 Product Size: 4 FL OZ (118 mL)
 Product Weight: 118 g

Artwork Type: IDM Commercial

Product Name: Mucinex Children's
 Product Description: Multi-Symptom Cold
 Product Type: Liquid
 Product Size: 4 FL OZ (118 mL)
 Product Weight: 118 g

Color Management

Color Management: CMYK
 Color Management: CMYK
 Color Management: CMYK

Barcode Info

Barcode Type: 1D
 Barcode Length: 12
 Barcode Width: 12
 Barcode Height: 12
 Barcode Color: Black

Health & Hygiene - Home

Health & Hygiene - Home
 Health & Hygiene - Home
 Health & Hygiene - Home

Product Size & Weight (Non-Automated)

Product Size & Weight (Non-Automated)
 Product Size & Weight (Non-Automated)
 Product Size & Weight (Non-Automated)

Product Size & Weight (Automated)

Product Size & Weight (Automated)
 Product Size & Weight (Automated)
 Product Size & Weight (Automated)

Product Size & Weight (Automated)

Product Size & Weight (Automated)
 Product Size & Weight (Automated)
 Product Size & Weight (Automated)

Product Size & Weight (Automated)

Product Size & Weight (Automated)
 Product Size & Weight (Automated)
 Product Size & Weight (Automated)

Product Size & Weight (Automated)

Product Size & Weight (Automated)
 Product Size & Weight (Automated)
 Product Size & Weight (Automated)



9212913

Drug Facts**Active ingredients**
(in each 5 mL)Guaifenesin 100 mg.....Expectorant
Phenylephrine HCl 2.5 mg.....Nasal decongestant**Purposes****Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

- Ask a doctor before use if the child has
- heart disease
 - high blood pressure
 - thyroid disease
 - diabetes
 - cough that occurs with too much phlegm (mucus)
 - persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

- Stop use and ask a doctor if
- nervousness, dizziness, or sleeplessness occur
 - symptoms do not get better within 7 days or occur with fever
 - cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl galate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

*as an OTC brand in the
Children's Cough/Cold category
(AlphaImpactRxProVoice Survey)

Other convenient forms for
Chest Congestion Relief



mini-melts™

On-the-go
granule packets for
chest congestion relief!

Keep carton for full information

Tamper evident: Do not use if neckband
on bottle cap is broken or missing.

Mucinex
Children's

**STUFFY NOSE &
CHEST CONGESTION**

Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

Pediatrician Recommended!

NDC 63824-951-07

Mucinex
Children's

**STUFFY NOSE &
CHEST CONGESTION**

Guaifenesin 100 mg - Expectorant
Phenylephrine HCl 2.5 mg - Nasal Decongestant

- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Alcohol-free



Ages 4+ yrs

**Very Berry
Flavor Liquid**

4 FL OZ (118 mL)

Mucinex
Children's

**STUFFY NOSE &
CHEST CONGESTION**

Each dose of Very Berry flavored
Mucinex Children's Stuffy Nose
& Chest Congestion Liquid:

- ✓ Relieves Stuffy Nose
- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- ✓ Alcohol-free



Dosing Cup Included

Questions or comments?
1-866-Mucinex (1-866-682-4639)
or www.mucinex.com

Dist by: RB Health (US)
Parsippany, NJ 07054-0224
©2020 RB Health
Made in Mexico
012221 3183776


HEALTH • HYGIENE • HOME

3163776



3 63824 27764 6

LOT CODE
EXP. DATE



8341440

112818

Drug Facts

Active ingredients Purposes (in each caplet)

Mucinex® FAST-MAX® DAY TIME SEVERE CONGESTION & COUGH

Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Active ingredients Purposes (in each caplet)

Mucinex FAST-MAX NIGHT TIME COLD & FLU

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Drug Facts (continued)

- temporarily relieves these common cold and flu symptoms (**NIGHT TIME only**):
 - cough
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
 - temporarily reduces fever (**NIGHT TIME only**)
 - controls cough to help you get to sleep

Warnings

- Liver warning (NIGHT TIME only):** This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 12 caplets in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. (**NIGHT TIME only**)
- with any other product containing diphenhydramine, even one used on the skin (**NIGHT TIME only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin (**NIGHT TIME only**)
- taking sedatives or tranquilizers (**NIGHT TIME only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**NIGHT TIME only**)
- marked drowsiness may occur (**NIGHT TIME only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT TIME only**)

USES

- temporarily relieves (**DAY TIME only**):
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - nasal congestion due to a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**DAY TIME only**)

Allergy alert (NIGHT TIME only):

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (NIGHT TIME only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Ask a doctor before use if you have

- liver disease (**NIGHT TIME only**)
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT TIME only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT TIME only**)

- avoid alcoholic drinks (**NIGHT TIME only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT TIME only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever (**DAY TIME only**)

Drug Facts (continued)

- pain, nasal congestion, or cough gets worse or lasts more than 7 days (**NIGHT TIME only**)
- fever gets worse or lasts more than 3 days (**NIGHT TIME only**)
- redness or swelling is present (**NIGHT TIME only**)
- new symptoms occur (**NIGHT TIME only**)
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Overdose warning (NIGHT TIME only): Taking more than the recommended dose (overdose) may cause liver damage. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period

Drug Facts (continued)

- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

**Inactive ingredients
(Mucinex FAST-MAX
DAY TIME SEVERE
CONGESTION & COUGH)**

corn starch, FD&C blue no. 2
aluminum lake, FD&C red no. 40
aluminum lake, magnesium stearate,
maltodextrin, mica, microcrystalline
cellulose, polyethylene glycol,
polyorbate 80, polyvinyl alcohol,
povidone, silicon dioxide, sodium
starch glycolate, stearic acid, talc,
titanium dioxide

**Inactive ingredients
(Mucinex FAST-MAX
NIGHT TIME COLD & FLU)**

corn starch, croscarmellose sodium,
crospovidone, FD&C blue no. 1
aluminum lake, FD&C blue no. 2
aluminum lake, ferric oxide yellow,
magnesium stearate, methacrylic
acid-ethyl acrylate copolymer (1:1)

Drug Facts (continued)

type A, mica, microcrystalline cellulose,
polyethylene glycol, polyvinyl alcohol,
povidone, silicon dioxide, sodium
bicarbonate, stearic acid, talc, titanium
dioxide

**PEEL CORNER TO READ
COMPLETE DRUG FACTS
AND INFORMATION**

Do not take the Mucinex Fast-Max DAY TIME SEVERE CONGESTION & COUGH and Mucinex Fast-Max NIGHT TIME COLD & FLU caplets at the same time.

Always wait at least 4 hours before taking another dose of Mucinex caplets.

Do not take more than a total of 12 caplets in any 24-hour period.

Take only as directed.

8337090

110718





8335593

060618

Keep outer package for
full information.

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects
to this phone number.

Maximum Strength per 4-hour dose
*Helps to relieve these symptoms
day or night

LIFT HERE TO OPEN

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Tamper evident: do not use if outer package
is opened or if blister is torn or broken.

www.mucinex.com

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches and pains
 - sore throat
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients corn starch, croscopovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

8335600

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224 ©2018 RB 060618

102519



**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Maximum Strength per 4-hour dose
Tamper evident: Do not use if neckband
on bottle cap is broken or missing.
*Helps to relieve these symptoms day or night

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Drug Facts

Active Ingredients **Purposes**
(in each 20 mL)

Acetaminophen 650 mg
.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....
.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....
.....Nasal decongestant

Drug Facts (continued)

Warnings

Liver warning: This product
contains acetaminophen.
Severe liver damage may occur
if you take:

- more than 6 doses in
24 hours, which is the
maximum daily amount
- with other drugs containing
acetaminophen
- 3 or more alcoholic drinks
daily while using this product

Allergy alert: Acetaminophen

2963



1201201

LOT:

EXP:

3186230

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).

Drug Facts (continued)

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)**Directions**

- do not take more than directed (see **Overdose warning**)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains:
 - sodium 9 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, D&C yellow no.10, edetate disodium, FD&C blue no.1, flavors, glycerin, propyl gallate, propylene glycol, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Drug Facts (continued)**Questions?**

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number

**COOLING
MENTHOL
FLAVOR**



Drug Facts

Active ingredients (in each liquid gel)

Purposes

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - stuffy nose
 - minor aches and pains
 - sore throat
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure

Drug Facts (continued)

- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



COLD & FLU

"Helps to relieve these symptoms day or night"
Maximum Strength per 4-hour dose
Do not take more than a total of 12 liquid gels in any 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

071817 3051674

MAXIMUM STRENGTH*

NDC 63824-015-66

Mucinex[®]
FAST-MAX[®]

**COLD, FLU
& SORE THROAT**

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant • **Phenylephrine HCl** – Nasal Decongestant

- ✓ Controls Cough, Thins & Loosens Mucus
- ✓ Nasal & Chest Congestion
- ✓ Sinus Pressure & Congestion
- ✓ Body Pain, Headache, Fever & Sore Throat

6 FL OZ (180 mL)

FOR AGES 12+

**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

***Per 4-hour dose**

Tamper evident: Do not use if neckband
on bottle cap is broken or missing.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org


3 63824 01565 1

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 20 mg.....Cough suppressant	
Guaifenesin 400 mg.....Expectorant	
Phenylephrine HCl 10 mg.....Nasal decongestant	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
 - nasal congestion
 - sore throat
 - stuffy nose
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224
©2017 RB
082317

LOT: 3051679
EXP:
MADE IN:

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
 - adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
 - children under 12 years of age: do not use

Drug Facts (continued)

Other information

- each 20 mL contains: sodium 8 mg
- store between 20-25°C (68-77°F)

9
SYMPTOM

■ DO NOT refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH ▸ HYGIENE ▸ HOME

Please visit our website
www.mucinex.com

RELIEF
3045695






8335592

061518

www.mucinex.com
Patents: www.rb.com/patents

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects
to this phone number.



LIFT HERE TO OPEN

**Maximum Strength
per 4-hour dose**
Keep outer package for
complete product information.



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Tamper evident: Do not use if outer package is opened or if blister is torn or broken.

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Uses temporarily relieves these common cold and flu symptoms:
■ nasal congestion
■ cough ■ minor aches and pains
■ sore throat ■ headache
■ temporarily reduces fever
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 12 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks daily while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have
■ liver disease ■ heart disease
■ diabetes ■ high blood pressure ■ thyroid disease
■ trouble urinating due to an enlarged prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see Overdose warning)
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years of age and older:
take 2 caplets every 4 hours
■ children under 12 years of age: do not use

Other information

■ store at 20-25°C (68-77°F)

Inactive ingredients corn starch, croscavidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

8357048

Dist. by: RB Health (US), Parsippany, NJ 07054-0224 ©2019 RB Health

041219





Drug Facts

Active ingredients (in each liquid gel)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches and pains
 - sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



Drug Facts

Active Ingredients Purposes (in each 20 mL)

Acetaminophen 650 mg
.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg
.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg
.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - headache
 - sore throat
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription

Drug Facts (continued)

or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children

Drug Facts (continued)

even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store between 20-25°C (68-77°F)
- dosing cup provided
- do not refrigerate

Inactive ingredients

anhydrous citric acid, D&C yellow no.10, edetate disodium, FD&C blue no.1, flavors,

PEEL CORNER TO REVEAL COMPLETE DRUG FACTS AND INFORMATION

*Per 4-hour dose
Tamper evident: Do not use if neckband on bottle cap is broken or missing.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3

LOT:

EXP:

3027764

Drug Facts (continued)

glycerin, propyl gallate, propylene glycol, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Please visit our web site
www.mucinex.com

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224

Made in England ©2016 RB



Drug Facts Active ingredients (in each liquid gel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Guafenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant		Purposes ■ temporarily relieves these common cold and flu symptoms: ■ headache ■ nasal congestion ■ minor aches and pains ■ cough ■ sore throat ■ temporarily reduces fever ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive		Drug Facts (continued) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin When using this product do not use more than directed Stop use and ask a doctor if ■ nervousness, dizziness, or sleepiness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 12 liquid gels in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 liquid gels in any 24-hour period ■ adults and children 12 years of age and over: take 2 liquid gels every 4 hours ■ children under 12 years of age: do not use			
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.		Other information ■ store at 20-25°C (68-77°F) ■ avoid excessive heat			
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland		Inactive ingredients FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water			
Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.					



3090468

012919


MAXIMUM STRENGTH

MULTI-SYMPTOM



www.mucinex.com
Patents: www.rb.com/patents

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects
to this phone number.



LIFT HERE TO OPEN

**Maximum Strength
Per 4-hour dose**
Keep outer package for
full information.



**Tamper evident: Do not use if outer package
is opened or if blister is torn or broken.**

Drug Facts

Active ingredients (in each caplet) Purposes

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with

Acetaminophen 325 mg ... Pain reliever/fever reducer
 Guaifenesin 200 mg ... Expectorant
 Phenylephrine HCl 5 mg ... Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - fever
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

- smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

3090854

Dist. by: RB Health (US), Parsippany, NJ 07054-0224 Made in England ©2019 RB Health 013019